



Article title: GLOBAL EVOLUTION AND PALEOGEOGRAPHIC DISTRIBUTION OF
MID - CRETACEOUS ORBITOLINIDS

Authors: Marcelle Boudagher-Fadel, G. David Price

License information: This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY) 4.0 <https://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Preprint statement: This article is a preprint and has not been peer-reviewed, under consideration and submitted to UCL Open: Environment for open peer review.

Links to data: Links to data:

||<http://www.ucl.ac.uk/ucl-press/browse-books/evolution-and-geological-significance-of-larger-benthic-foraminifera>

DOI: 10.14324/111.444/000013.v3

Publication date: 19 March 2019

Keywords: Foraminifera, Orbitolinids, , mid-Cretaceous,, biostratigraphy,, phylogeny,, palaeogeographic distribution,, Extinctions, Global anoxic events, Sea-level changes, The Environment, Climate, Ecology

GLOBAL EVOLUTION AND PALEOGEOGRAPHIC DISTRIBUTION OF MID - CRETACEOUS ORBITOLINIDS

MARCELLE K. BOUDAGHER-FADEL* AND G. DAVID PRICE

Department of Earth Sciences, UCL, Gower Street, London, WC1E 6BT, UK.

***m.fadel@ucl.ac.uk**

ABSTRACT

Members of the Larger Benthic Foraminiferal (LBF) family Orbitolinidae occurred from the Cretaceous to the Paleogene, however, they were most diverse during the mid-Cretaceous, and dominated the agglutinated LBF assemblages described from limestones of that period. Various orbitolinid species have been used to zone and date lithologies formed in the shallow, warm waters of the Aptian to the Early Cenomanian, and many, sometimes inaccurate, generic and sub-generic nomenclatures have been proposed to differentiate the often subtle morphological changes that orbitolinids exhibit over time. Until now, it has not been possible to develop an effective global overview of their evolution and environmental development because descriptions of specimens from Asia have been relatively rare. Following our recent study of over 1800 orbitolinid-rich thin sections of material from 13 outcrops of the Langshan Limestone, from the Southern Tibetan Plateau, and from the Barito Basin, South Kalimantan, Indonesia, it has been possible to compare the stratigraphic ranges of these orbitolinids with previously described Tethyan and American forms, based on the use of a planktonic zonal (PZ) scheme, itself tied to the most recent chronostratigraphic scale. This has allowed the reconstruction of the phylogenetic and paleogeographic evolution of the orbitolinids from their Valanginian origin in the Tethys. Although Tethys remained the paleogeographic focus for the orbitolinids, it is inferred here for the first time that a bi-directional paleogeographic migration of some orbitolinid genera occurred from Tethys to the Americas and also to the Western Pacific region. Our observations and dating confirm that global marine regressions in the Aptian were coincident with, and may well have facilitated, these orbitolinid transoceanic migrations. Migration stopped, however, after rising sea-level in the Early Albian appears to

have again isolated these provinces from each other. Tectonic forces associated with the subduction of the Farallon Plate and further sea-level raises led to the opening of the Western Interior Seaway in the North America, which correlates with, and may have been the cause of, the Middle Albian (top of PZ Albian 2) extinction of the American orbitolinids. The extinction of the orbitolinids revealed that the Western Pacific province was split into two sub-provinces, with extinction occurring at the end of the Early Albian (top of PZ Albian 1) in the Northwest Pacific sub-province, and at the end of the Albian (top of PZ Albian 4) in the sub-province that is today South East Asia (on the margins and west of the Wallace Line). The final virtual extinction of the orbitolinids occurred at the end of the Cenomanian in the Tethyan province, which coincides with, and may have been caused by, global anoxic oceanic events that correlate with a near-peak Mesozoic eustatic sea-level high-stand that led to the overall global collapse of the paleotropical reef ecosystem at that time.

INTRODUCTION

The Orbitolinidae are an agglutinated, but now extinct, family of the Larger Benthic Foraminifera (LBF). Orbitolinidae were present in the warm, shallow marine waters of the Early Cretaceous to the Early Oligocene, however, they were most diverse during the mid-Cretaceous. During the Early to mid-Cretaceous (Valanginian to Early Cenomanian), there was an identifiable increase in the complexity of their morphological structure, which enabled them to house within their tests symbiotic algae (BouDagher-Fadel, 2018a), and it is these forms which are the subject of this paper. Traditionally, orbitolinids are considered to define two major, distinct paleogeographic realms, namely those of the Americas and the Tethys (see BouDagher-Fadel, 2018a), but in this study we find forms reported from the Western Pacific are distinct from their Tethyan forebears, and so define a third orbitolinid province.

The symbiotic orbitolinids were reef-forming organisms, and they are found in association with other marine forms, including planktonic foraminifera. This coexistence with planktonic forms, enables their stratigraphic ranges to be defined very precisely, as they can be tied to the high resolution planktonic zonal (PZ) dating scheme of BouDagher-Fadel (2018b; see Fig. 1), which itself is tied to the absolute time scale of Gradstein et al. (2012).

Early to mid-Cretaceous orbitolinids have been described from ancient Tethyan limestones from, for example, the Mediterranean (Husinec, 2001), Southwest England (Carter and Hart, 1977; Hart et al., 1979; Hart, 1982; Hart and Williams, 1990; Simmons et al., 2000; Simmons et al., 1992), Spain (Vilas et al., 1995; Caus et al., 1997), Israel, Lebanon and Syria (Saint-Marc, 1970), Yemen (Cherchi et al., 1998), Oman (Simmons and Hart, 1987; Simmons, 1994; Masse et al., 1998), Saudi Arabia (Hughes, 2005; Cantrell et al., 2017), the United Arab Emirates (Vahrenkamp, 1996), Iran (Mehrnusch, 1973; Shakib, 1994; Roozbahani, 2011), Afghanistan (Schroeder, 1975), and Tibet (Zhang, 1982, 1986; BouDagher-Fadel et al., 2017). They are also reported from the Northwest Pacific (Iba et al., 2011), Japan and Sakhalin (Yabe and Hanzawa, 1926; Hofker, 1963; Ujiie and Kusukawa 1968; Matsumaru 1971, 1973; Matsumaru et al. 1976; Salnikov and Tikhomolov 1987; Iba et al., 2005; Matsumaru, 2005; Matsumaru and Furusawa 2005, 2007; Iba and Sano 2006; Matsumaru and Furusawa 2007), and in Africa, where they are found in Ethiopia (Bosellini et al., 1999), Somalia (e.g. Luger et al., 1990; Bosellini, 1992; Cherchi and Schroeder, 1999) and Tanzania (Peybernès, 1982).

Furthermore, orbitolinids have been reported from the northwestern Atlantic, off the Flemish Cap, Newfoundland (Sen Gupta and Grant, 1971; Schroeder and Cherchi, 1979), and have been described from the Caribbean and the Americas (Douglass 1960), Mexico (Meza, 1980; Pantoja-Alor et al., 1994; Omaña and Alencáster, 2009), and Venezuela (Görög and Arnaud-Vanneau, 1996).

During their existence, the structurally complex orbitolinids showed relatively rapid phylogenetic evolution, developing many stratigraphically short-ranged species, which when combined with the PZ scheme (see Fig. 1) act as a very important and precise index fossil group for the shallow-marine environments of the mid-Cretaceous Tethys (Simmons et al., 2000; Schroeder et al., 2010; BouDagher-Fadel et al., 2017; BouDagher-Fadel, 2018a). As a result, they have been widely adopted as a biostratigraphic tool by industry in the exploration of Middle Eastern and other oil and gas fields.

In this paper, the evolution and paleogeographic development of these symbiotic, morphologically complex orbitolinids is inferred from the re-analysis of the published data referred to above, and combined with new observations from over 1800 thin sections of material from 13 outcrops of the Langshan Limestone of the Southern Tibetan Plateau (see Fig. 2), the Sangzugang Formation in Southern Lhasa subterrane (see An et al., 2014), the Xiagezi-II section of the Langshan Formation in southern part of Northern Lhasa subterrane (see Sun et al., 2015), the Azhang and Guolong sections from the Langshan Formation in Northern Lhasa subterrane (see BouDagher-Fadel et al., 2017), the Jingshughan, Langshan, Xiongba, Xiongmei, Baoji, Daya, Gegi, Letie and Zulong sections (Yiwei et al., in preparation), and the Jiarong and Laxue sections from the Linzhou Basin (see BouDagher-Fadel et al., 2017). In addition, material has been studied from the western flank of the Meratus Mountains, an uplifted accretionary collision complex that records suturing of East Java-West Sulawesi to the Sundaland margin during the mid-Cretaceous (see Fig. 3). The uplifted complex now forms the eastern boundary of the Barito Basin, South Kalimantan, Indonesia (see Witts, 2011).

By correlating these observations and literature data with our high resolution PZ scheme (BouDagher-Fadel, 2018b, Fig. 1), we are able to infer, for the first time, a comprehensive, global synthesis of the biostratigraphic, phylogenetic and paleogeographic evolution of these orbitolinids. We infer that the earliest morphologically complex orbitolinids evolved in the Tethyan from primitive Valanginian forms such as *Valdanchella*, *Paleodictyoconus*, and *Campanellula* (Fig. 4). More complex forms developed rapidly into different Tethyan

phylogenetic lineages (e.g. Figs 4 and 5). It appears that major, global sea-level regressions starting in the Early Aptian (PZ Aptian 1, 125.0 Ma) and in the Late Aptian (PZ Aptian 4, 116.5Ma; see Fig. 6), correlate with and probably facilitated bi-directional transoceanic migration of orbitolinids. One migration was from the Tethyan to the previously recognised American province, but a second migration was from Tethys to the newly defined Western Pacific province (see Fig 7). These migrations stopped after rising sea-level during the Early Albian (PZ Albian 1) appears to have isolated the provinces one from another.

The isolated orbitolinids of the Northwest sub-province of the Western Pacific (present day Japan) became extinct at end of the Early Albian (top of PZ Albian 1), while those in the isolated American province became extinct at the end of PZ Albian 2 (106.7Ma). All forms in the sub-province that is today South East Asia (on the margins and to the west of the Wallace Line) went extinct at the end of the Albian (top of PZ Albian 4. The “hotspot” for orbitolinid, evolution, however, remained in Tethys, where environmental conditions continued to contribute to their success until the end of the Cenomanian, when virtually all symbiotic, morphologically complex orbitolinids became extinct, as indeed did many of the other agglutinated LBF that dated from the Early Cretaceous and Jurassic (see BouDagher-Fadel, 2018a). These extinctions coincided with an anoxic oceanic event (Bambach, 2006), and correlate with a near-peak Mesozoic eustatic sea-level high-stand (see Fig. 6, and Miller et al, 2005).

MORPHOLOGICAL CHARACTERISTICS OF ORBITOLINIDS

The orbitolinids are members of the order Textulariida, which have agglutinated tests that are made of foreign particles bound by organic cement. They are characterized by having conical tests, subdivided into numerous chambers, and are usually a few millimetres in height and diameter (although as noted, some forms attained diameters of 5 cm or more). The numerous uniserial discoidal chambers are partially subdivided by radial or transverse partitions, or pillars. They have cribrate, areal apertures (see Fig. 8).

The Cretaceous morphologically complex orbitolinids are divided into the dictyoconines and orbitolinines, and range from the Valanginian to the Cenomanian. They are divided into the following five morphological groups, (see BouDagher-Fadel, 2018a):

- (i) Orbitolinids with no complex central zones (e.g., *Campanellula*, PZ Valanginian 1). They lack thick radial partitions and pillars in the central zone.

- (ii) Orbitolinids with a complex central zone and radial partitions thickening away from the periphery and breaking up into pillars in the central zone, first appeared in the Late Valanginian with developed peripheral tiered rectangular chamberlets. They evolved into the dictyoconines (e.g., *Paleodictyoconus*, PZ Valanginian 2, Fig. 4; *Paracoskinolina*, PZ Barremian 1-Albian 4), or into the orbitolinines (e.g., *Urgonina*, PZ Barremian 1, Fig. 4) from forms with the outer parts of their chambers lacking partitions but with interseptal pillars connecting the adjacent septa.
- (iii) Orbitolinids with radial partitions thickening away from the periphery to anastomose centrally around the aperture and form a reticulate zone in transverse section, also first appeared in the Late Valanginian (e.g., *Valdanchella*, PZ Valanginian 2). The peripheral zones of their chambers are subdivided into rectangular chamberlets by fine radial partitions (Fig. 4).
- (iv) Orbitolinids with radial partitions that became zigzag, thickening and fusing centrally, giving a stellate appearance in transverse section, first appeared in the Aptian (e.g., *Simplorbitolina*, PZ Aptian 1). Their tests may have tiered peripheral chamberlets (e.g. *Dictyoconella*, PZ Cenomanian 3, Fig. 4).
- (v) Orbitolinids with radial partitions thickening, with triangular cross-sections away from the periphery and anastomosing in the central area, first appeared in the Barremian (e.g. *Eopalorbitolina*, PZ Barremian 1, Fig. 5) and evolved rapidly in the mid-Cretaceous. The test of these orbitolinids is defined by the shape of the embryonic apparatus, and by the size and shape of the chamber passages that can be seen in tangential sections. The earliest formed chambers of the megalospheric generation can form a complex embryonic apparatus, which can be divided into a protoconch, a deuteroconch, a sub-embryonic zone and peri-embryonic chamberlets (see Plate 1, figs 2, 5; Fig. 4). In axial section, the embryo is located at the apex of the cone, followed by a series of discoidal chamber layers. The embryonic apparatus evolved from a simple apparatus, consisting of a large globular fused protoconch and deuteroconch, followed by peri-embryonic chambers as in *Palorbitolina*, to an embryonic apparatus divided into a protoconch and deuteroconch but a not completely divided sub-embryonic zone, as in *Praeorbitolina*. This latter evolved in turn into forms in which the deuteroconch and sub-embryonic zone are more or less of equal thickness, as in *Mesorbitolina* (Plate 1, fig. 4). In *Conicorbitolina* (Plate 3, fig. 6) the marginal zone became extensively divided by vertical and

horizontal partitions, while in *Orbitolina* the deuteroconch is highly subdivided and of much greater thickness than the sub-embryonic zone (see Figs 4, 5; Schroeder, 1975; Hottinger, 1978; Simmons et al., 2000; BouDagher-Fadel, 2008; Schroeder et al., 2010; BouDagher-Fadel et al., 2017). In transverse section, the chambers are seen divided into a marginal zone, with sub-epidermal partitions, and a central zone with radial partitions (Plate 1, fig. 1; Figs 8.1-8.2, 9). The chamber passages are formed in the radial part of the central zone of each chamber layer (Figs 8.1-8.2, 9h), where each chamber passage is subdivided by vertical main partitions, which are prolongations of the vertical main partitions of the marginal zone (Figs 9a-f, h). The radial partitions (Fig. 9f) in advanced orbitolinids (e.g. *Mesorbitolina*, *Orbitolina*) thicken away from the periphery and anastomose in the central area, producing an irregular reticular network (Plate 1, figs 3, 6-8; Figs 9g, i-j; Fig. 10b). In cross section, the chamber passages can be triangular (Figs 9c, 10a), rectangular (Fig. 10c) or oval, or can show a gradation between shapes (Fig. 9e) (Schroeder, 1975). In the radial zone of *Orbitolina*, the stolons are arranged in radial rows alternating from one chamber to the next one (see BouDagher-Fadel, 2018a). Their alternating position would have obliged the protoplasm to flow in an oblique direction (Hottinger, 1978). In the annular radial zone of the conical test (Plate 1, figs 1, 3, 7), radial septula subdivide the chambers into radial compartments with various thickness and textures (Plate 1, fig. 8; Fig. 9f, h, k-r), narrowing towards the centre to fuse into a reticular network (Plate 1, figs 1, 6, 8; Fig. 9g, i, j) which minimizes the volume of chamberlet cavities (Plate 1, figs 1, 6-8).

BIOSTRATIGRAPHY, PHYLOGENY AND PALEOGEOGRAPHIC DISTRIBUTION OF THE ORBITOLINIDS

The orbitolinids are very useful biostratigraphic markers in early to mid-Cretaceous Tethyan carbonate platforms (Henson, 1948; Schroeder, 1975; BouDagher-Fadel et al., 2017). They have short ranges and are, with practice, easily identified in thin sections (e.g. see Plates 2 and 3). Orbitolinids, unlike some LBFs of the period (e.g. the miliolides), show provincialism. Traditionally, they are considered to define two major, distinct paleogeographic realms, namely those of the Americas and the Tethys (see BouDagher-Fadel, 2018a).

Many forms from the morphological Group (i) described above evolved gradually to more advanced forms from Groups (ii) to Groups (vi). Notable and characteristic lineages include:

- *Campanellula* – *Paracoskinolina* – *Coskinolinoides* – *Dictyoconus* (PZ Valanginian 1-Cenomanian 3; all forms in this group became extinct in the Cenomanian, except *Dictyoconus*, which persisted to the Oligocene). The evolutionary trend of this Group (i) – Group (ii) lineage is characterized by an increase in test diameter and the development of increasingly complex radial partitions radial partitions thickening away from the periphery, that break up into pillars in the central zone, forming highly developed and complex layers of chamberlets.
- *Valdanchella* – *Paleodictyoconus* – *Montseciella* – *Rectodictyoconus* – *Simplorbitolina* – *Neorbitolinopsis* (PZ Valanginian 1-Cenomanian 1). The evolutionary trend of this Group (i) to Group (v) lineage is characterized by the increase in size, a gradual enlargement of the whole embryo the development of the megalospheric embryo in a centric or near centric position, and the development of increasingly complex radial partitions, becoming zigzag with a stellate appearance in transverse section (as in *Simplorbitolina*) or thickened and fused centrally forming highly developed and complex layers of chamberlets (as in *Neorbitolinopsis*).
- *Urgonina* – *Eopalorbitolina* – *Palorbitolina* – *Palorbitolinoides* (PZ Barremian 1-Albian 1). The evolutionary trend of this Group (ii) – Group (v) lineage (see Fig. 5) is related to the formation of peripheral, tiered, rectangular chamberlets in two or more series, the shape and position of the embryonic apparatus from a bi-chambered embryo in a clear eccentric position, with a missing peri-embryonic zone in *Eopalorbitolina* (see Fig. 5), to the development and the increase in size of the peri-embryonic zone to embrace more and more of the embryonic chamber, surrounding the upper half of the nearly centric embryonic chamber in *E. transiens*, and becoming completely annular surrounding the upper part of the centric embryonic chamber in *Palorbitolina lenticularis* (Figs 5, 10). In *Palorbitolinoides* (e.g. *P. hedini*, Plate 2, fig. 5) the large and flattened embryonic chamber is surrounded by a developed inflated peri-embryonic zone.
- *Praeorbitolina* – *Mesorbitolina* – *Orbitolina* – *Conicorbitolina* (PZ Aptian 1-Cenomanian 3). The main evolutionary characters in this Group (v) lineage are the position of the embryonic apparatus, which is in an eccentric position in earlier forms (e.g. *Praeorbitolina*), but centrally placed in advanced forms, consisting of the protoconch and the deuterconch (e.g. *Mesorbitolina*). In *Conicorbitolina* (PZ Albian 4-Cenomanian 1) the

large proloculus is divided into a protoconch and deuteroconch, with the marginal zone becoming extensively divided by vertical and horizontal partitions (Plate 3, fig. 6; Fig. 5). The main evolutionary characters of this *Orbitolina* (PZ Albian 3- Cenomanian 3) is the increase in size of the apically situated embryonic apparatus, where the deuteroconch becomes about three times thicker than the sub- embryonic zone (Figs 5, 9).

On the basis of this study, and using the lineages described above, we are able to establish for the first time that there were in fact three distinct paleogeographic provinces for these symbiotic, morphologically complex orbitolinids (see Fig. 7); namely the previously defined American province (including current day Texas, Venezuela, Mexico), a Tethyan province (including Europe and the Flemish Cap off Newfoundland, Arabia, Turkey, Iran, Lebanon, Oman, Syria, Qatar, Tibet), and a newly identified Western Pacific province, which is divided into two sub-provinces; the sub-province of Northwest Pacific, which include Japan and the Philippine island of Cebu, and a sub-province that includes what is today South East Asia (west of the Wallace Line).

In Tethys, morphologically complex orbitolinids and their precursors are common from the Valanginian (PZ Valanginian 1) to the Cenomanian (PZ Cenomanian 3), and exhibit several of the phylogenetic lineages described above, while in the Americans orbitolinids are only found between the Early Aptian (PZ Aptian 1) and Middle Albian (PZ Albian 2), and are predominantly represented by the Group (v) genera *Palorbitolina* and *Mesorbitolina*. In the Western Pacific, unidentified and unconfirmed orbitolinids have been listed in the literature as dating from the Late Hauterivian to the Early Aptian (see Iba et al., 2011). These early forms are however contested, but the Group (v) *Praeorbitolina* – *Mesorbitolina* lineage is definitely confirmed from PZ Aptian 1 to PZ Albian 1 in Northwest Pacific sub-province and to PZ Albian 4 in South East Asia sub-province.

From this global pattern, we infer that the original hotspot for the evolution of the complex orbitolinids was Tethys, but as will be described below, following migration events in the Early Aptian out of Tethys, some lineages of the orbitolinids spread to the other provinces. It seems that the migration stopped after the Early Albian, and that the provinces were again isolated. There then developed provincial, parallel, but specifically distinct evolutionary trends, until the subsequent provincial extinctions in the Americas and the West Pacific (see Fig. 11).

The Tethys

Throughout the Tethyan province orbitolinids of Groups (i-v) evolved many lineages. They became morphologically complex and widespread, and are often associated with calcareous algae. Their principle Tethyan lineages which evolved from Group i include:

- Group (ii) – Group (iv) *Valdanchella* – *Paleodictyoconus* – *Montseciella* – *Rectodictyoconus* – *Simplorbitolina* – *Orbitolinopsis* (PZ Valanginian 1-Cenomanian 1);
- Group (v): *Eopalorbitolina* – *Palorbitolina* – *Palorbitolinoides* (PZ Barremian 1-Albian 1);
- Group (v): *Praeorbitolina* – *Conicorbitolina* (PZ Aptian 1-Cenomanian 1).

The orbitolinid associations of western Tethys, southern Neo-Tethys margin and of southwest Europe are similar to those of the Tibetan carbonate platforms, and they all form a part of the Tethyan realm (BouDagher-Fadel et al., 2017). All cosmopolitan orbitolinids appeared in Tethys before spreading to other provinces. For example, in Tethys, *Palorbitolina lenticularis* (Plate 2, fig. 1; Fig. 10) first occurred in Late Barremian (PZ Barremian 3, 127 Ma; BouDagher-Fadel et al., 2017), 2 million years before its first appearance in the American and Western Pacific provinces at the beginning of Aptian (PZ Aptian 1, 125.0 Ma). The oldest *P. lenticularis* recorded in what is today the “American” continent was recorded by Schroeder and Cherchi (1979) from the Late Barremian of the Flemish Cap, North West Atlantic. From the palaeogeography of the time, however, we infer that at this stage the Flemish Cap was the extreme extension of the north western Tethyan realm and was isolated from the more southerly parts of the American province (see Fig. 7).

The earliest *Mesorbitolina* (e.g., *Mesorbitolina lotzei*) appeared first in Tethys, in PZ Aptian 2. The cosmopolitan *M. parva* – *M. texana* first appearing in the Late Aptian (PZ Aptian 3, 119.5Ma), 3 million years earlier than in the American and Western Pacific provinces where their first appearances are recorded in PZ Aptian 4, 116.5Ma (Douglas, 1960; Simmons et al., 2000; Iba et al., 2011; BouDagher-Fadel et al., 2017). Subsequent to its first appearance, the Tethyan *Mesorbitolina* evolved many phylogenetic lineages, which show the typical evolution from having a simple embryonic apparatus to developing a more complicated one. The most common Late Aptian – Cenomanian (PZ Aptian 4-Cenomanian 1) lineage being the *M. texana* – *M. aperta* lineage (see Plate 1, figs 1-3, 6; Plate 2, fig. 4; Fig. 5), where the open

deuteroconch in the square embryonic apparatus evolves into a deuteroconch subdivided in the upper part by several partitions of different sizes, whereas the lower part exhibits an irregular network of partitions (BouDagher-Fadel et al., 2017). No equivalent lineage is found in the other provinces, suggesting that by this stage the provinces were again isolated one from another.

All Tethyan orbitolinids became extinct (with the exception of *Dictyoconus*) at the end of the Cenomanian.

The Western Pacific

In the Western Pacific province, orbitolinids limestones, associated mainly with Cretaceous arc volcanics, form two sub-provinces. One occurs north along the Eurasian continental margin to the Philippines and Japan, and the other is to the south, along a belt near the Early Cretaceous margin of Sundaland, in what is today South East Asia (Hofker 1963; Hashimoto et al., 1975).

In the Northwest Pacific sub-province, reported occurrences of orbitolinids are patchy with numerous doubtful identifications, but those which are definitely identified belong to Group (v). *Palorbitolina lenticularis* is first recorded from the beginning of PZ Aptian 1 (125.0Ma, 2 million years after its first appearance in Tethys) in the eastern Philippines (Cebu) and Japan (Hashimoto et al. 1978; Iba et al., 2011).

In the South East Asian sub-province, orbitolinids are more common and occur from PZ Aptian 1 (125.0 Ma) to PZ Albian 4 (100.5 Ma). In West Sarawak and NW Kalimantan, orbitolinid-rich beds are recorded from the Early Aptian (PZ Aptian 1) of Pedawan and Seberoeang Formations (Hashimoto and Matsumaru, 1977). In North-Central Kalimantan orbitolinids are documented from the Aptian to Early Albian (PZ Aptian 1 – Albian 1) of the Selangkai Formation in the Upper Kapuas River region, (Hashimoto et al, 1975). Other Early Cretaceous orbitolinid localities include SE Kalimantan along the Meratus Mountains front East of Martapura (Hashimoto and Matsumaru, 1974), in South Sumatra, Ratai Bay, Lampung (Yabe, 1946), in the Gumai Mountains (Muspér, 1937), Central Java (Harloff 1933), and West Sulawesi (Brouwer, 1934, White et al., 2017).

The Western Pacific orbitolinids are mainly of Tethyan origin belonging to Group (v). Aptian forms originally described as endemic to the South East Asia sub-province are in fact found to be synonyms to the Tethyan forms. As an example, *Orbitolina scutum* and *O. trochus*, originally named as *Patellina scutum* and *P. trochus* are both described from Borneo and were

assumed to be of Eocene age by Von Fritsch (1879), but were later re-identified as the Tethyan species *Palorbitolina lenticularis* and *Mesorbitolina parva* (Hashimota and Matsumaru 1974). While *P. lenticularis* ranges from Late Barremian to Early Aptian in Tethys, it is only recorded from the Aptian in the Western Pacific.

In addition to these forms, we record in this work for the first time the presence of the Late Aptian to Early Albian (PZ Aptian 3 to Albian 1) Tethyan species of *Palorbitolinoides orbiculata* (Plate 2, fig. 3; Plate 3, figs 4, 7) in the Early Albian (PZ Albian 1) of the western flank of the Meratus Mountains, Barito Basin, Southeast Kalimantan, Indonesia. The Tethyan genus *Conicorbitolina* which evolved from *Mesorbitolina* in Tethys in Albian 3 and ranges to Cenomanian 1 (see Fig. 7) is also recorded here for the first time from the Late Albian (PZ Albian 4) of the Barito Basin, Kalimantan, *Conicorbitolina* sp. (Plate 3, fig. 6). Although the shape of the test is similar to the Tethyan *C. conica*, those of the Southeast Kalimantan have variations in the shape and number of perieymbryonic chambers (see Plate 3, fig. 6). This is an example of parallel evolution, which gave rise to a similar but distinct form from that found in the Tethyan province. We infer, therefore, that following their initial migration to the Western Pacific, the *Mesorbitolina* lineages subsequently exhibited parallel lineages evolving at different rates within the two provinces, with no further exchange of forms apparent, suggesting that the provinces again became isolated one from another.

The *Praeorbitolina* – *Mesorbitolina* lineages are represented in the Western Pacific province by *Praeorbitolina cormyi* (Plate 2, fig. 2), *P. wienandsi*, *Mesorbitolina parva*, and *M. texana* (Plate 1, figs 7-8; Plate 3, figs 1-3), and have been recorded from the Late Aptian to Early Albian (PZ Aptian 3 -Albian 1), again 5.5 million of years after their first appearance in Tethys. *Mesorbitolina subconca* (Plate, 1, fig.5; Plate 2, fig. 6; Plate 3, fig. 5) is recorded here for the first time from the Early Albian (PZ Albian 1) of the Barito Basin, Southeast Kalimantan, 3.5 million of years after its first appearance in Tethys.

Groups (i-iv) forms, unlike in the Tethys, seem to be missing from the Western Pacific province. Also, unlike the Tethyan realm, the orbitolinids do not survive the Albian-Cenomanian boundary, but disappeared completely from Japan at the end of PZ Albian 1 (Iba et al., 2011) and, as shown here, from the Barito Basin, Southeast Kalimantan, Indonesia at the end of PZ Albian 4. No orbitolinids are known from east of the Wallace Line in East Indonesia and Australia-New Guinea regions (Gorsel 2014), as these foraminifera required a tropical shallow marine settings, which was not present at this time along the North West Australian margin.

The Americas

Tethyan orbitolinids belonging to Groups (ii) and (v) seem to have migrated into the American province, however at a much later date than their first appearance in Tethys.

The American province, unlike the Western Pacific province, contains representatives of the dictyoconines from Group (ii). *Paracoskinolina*, which first appeared in the Barremian (PZ Barremian 1) of Tethys, and *Dictyoconus*, which first appeared in the Aptian (PZ Aptian 1) of Tethys, first appeared in the Albian (PZ Albian 1) of Texas, Mexico, and Venezuela (Mayne, 1955; Arnaud Vanneau and Sliter, 1995), and range to PZ Albian 2. Species such as *Paracoskinolina sunnilandensis* (PZ Albian 2) and *Dictyoconus walnutensis* (PZ Albian 1-2) are unique and indigenous to the American province, and forms recorded as the same as Tethyan species are in fact incorrectly defined. This unique occurrence excludes a West to East migration (Cherchi, 2004), and confirms that for large parts of the Albian the American and Tethyan provinces were ecologically isolated one from another.

The earliest form from Group (v) reported from the American province is *Palorbitolina lenticularis* from PZ Aptian 1 (125.0Ma) in deposits of south Mexico, and appear 2.0 million years later than its first occurrence in the Late Barremian of Tethys. The Tethyan *Mesorbitolina* are also widespread in the bank and reef deposits of Texas, New Mexico, Arizona, Guatemala, Honduras and Venezuela (PZ Aptian 4 - Albian 2). The cosmopolitan forms, *Mesorbitolina texana* - *M. parva* group occurring from PZ Aptian 3- Albian 1 (119.5-109.8Ma) in Tethys (BouDagher-Fadel et al., 2017), but are only reported from the PZ Aptian 4-Albian 2, (116.5-109.8Ma) of Texas, with *M. parva* only found in the PZ Albian 2 of the Americas.

In the Early Albian, species of *Mesorbitolina* continued to thrive in the Americas but developed provincial specific forms, not found in the Tethys or Western Pacific provinces. Thus, the American lineage *M. minuta* - *M. gracilis* - *M. crassa* of the PZ Albian 1-2 (Douglass, 1960; Monreal and Longoria, 1999) indicates that once the orbitolinids were established in the American province in the latest Aptian, they evolved independently from, yet in a parallel way to, their Tethyan ancestors, by means of gradual development of their embryonic apparatus. Those American species that had been previously reported from the Tethys or the Western Pacific were in fact mis-identified. For example, the American *Mesorbitolina minuta* was reported by Matsumaru and Furusawa (2007), from central Hokkaido, but was corrected as a synonym of *M. texana* by Cherchi and Schroeder (2009).

DISCUSSION

The Early Cretaceous is believed to have been a greenhouse period, with high atmospheric CO₂ concentrations (Royer et al., 2007), high global average temperatures with sea-surface temperatures exceeding 32°C (Skelton and Masse, 2000; Littler et al., 2011), and a stable climate (Littler et al., 2011). The earliest Cretaceous (Berriasian-Hauterivian) was also characterised by a sustained period of global low sea-levels, which were replaced in the Barremian by a significant global sea-level transgression (see Fig. 6), reaching its maximum at around 129 Ma, Barremian 2. This sea-level rise flooded low-lying continental regions and so created new ecological niches around the globe, one of which was filled in Tethys by the evolving orbitolinids.

The globally warm period continued in the mid-Cretaceous and was characterized by an increase in the number of agglutinated foraminiferal forms having large alveoles, such as the litiolid *Pseudocyclamina*, or forms with internal radial partitions, such as the orbitolinids (see BouDagher-Fadel, 2018a). This may have been an adaptation to the extreme climatic and oceanic conditions (increases in temperature and oceanic anoxia; e.g., Kerr, 2006) during this interval (BouDagher-Fadel, 2008), linked to an inferred dramatic increase of carbon dioxide in the atmosphere possibly triggered by enhanced global volcanism (e.g., the Ontong Java flood events). The high CO₂ levels during this greenhouse period also would have led to increased oceanic acidity (Naafs et al., 2016), which would have favored the ecological domination of the Textulariida, exemplified by the orbitolinids with their agglutinated tests, over those forms with biogenically precipitated calcitic tests that dominated before and after this period.

Evolving from earlier Valanginian forms, by the Late Barremian (PZ Barremian 3), major new lineages of the agglutinated orbitolinids had appeared in Tethys (see Cherchi and Schroeder, 2004). These robust forms had the ability to survive in many shallow carbonate environments (Arnaud-Vanneau, 1980), however, they were most common in the outer platform (Vilas et al., 1995; BouDagher-Fadel, 2008, 2018a; and see Fig. 12).

As noted above, we have shown that all cosmopolitan orbitolinids appeared in Tethys before migrating to other provinces. We have seen that once established in the American and Western Pacific provinces, local provincial forms evolved, indicating that they were subsequently again isolated from the Tethyan province. In previous studies of Cenozoic LBF,

specifically the lepidocyclinids (BouDagher-Fadel and Price, 2010), the miogypsinids (BouDagher-Fadel and Price, 2013), the nummulitoids (BouDagher-Fadel and Price, 2014) and the orthophragminids (BouDagher-Fadel and Price, 2017), we have observed similar developments, with periods of migration from one province to another followed by subsequent isolation and development of local provincial lineages. In these Cenozoic cases, the periods of inter-provincial migration coincided with major sea-level regressions, while the subsequent provincial isolation coincided with global sea-level transgressions. As observed in this study, it appears that a similar correlation occurs with the Cretaceous orbitolinids, with migrations from Tethys occurring during the time of Aptian sea-level low stands (Fig. 6), followed by isolation when the sea-level again rose in the Albian.

Thus, in Tethys, *Palorbitolina lenticularis* (Plate 2, fig. 1; Fig. 10) first occurred in Late Barremian (PZ Barremian 3, 127 Ma), 2 million years before its first appearance in the American and Western Pacific provinces (at the beginning of Aptian, PZ Aptian 1, 125.0 Ma). This migration coincides with the global sea-level regression that marks the beginning of PZ Aptian 1, and which culminates with the global sea-level minimum at the end of PZ Aptian 2.

Similarly, the earliest *Mesorbitolina* (e.g., *Mesorbitolina lotzei*) appeared first in Tethys, in PZ Aptian 2, but are not recorded until PZ Aptian 3 in the Western Pacific province, and PZ Aptian 4 in the American province.

After the earliest migration in the Aptian, the American Province appears to have been again isolated from Tethys throughout the later Albian and the more advanced lineages of Group (v) (e.g. *Orbitolina*, *Conicorbitolina*) of the Tethyan provinces, which appeared in Late Albian, are not found in the Americas. The evolutionary patterns inferred from Tethyan species diverge from those observed in the Americas, confirming that these two provinces were isolated from each other at this time. The progressive changes seen in the different lineages is regarded here as an example of orthogenesis, which resulted in the development of morphologically similar yet phylogenetically distinct forms with distinct biostratigraphic and paleogeographic characteristics.

The American orbitolinids became extinct at the end of the PZ Albian 2, 12.8 Ma earlier than those of Tethys (end Cenomanian 3). This event corresponds to the opening of the Western Interior Seaway triggered by sea-level rises and tectonic forces associated with the subduction of the Farallon Plate in the Late Albian. This produced for a period an epicontinental sea over

western North America that linked the tropical seas with a previously separate Arctic Ocean. This fully open seaway persisted in the Albian and the Cenomanian, flooding the orbitolinids habitats with cooler deeper waters, and was probably the cause of the orbitolinids extinction in the American province.

In the Western Pacific province, the Late Aptian to Early Albian larger benthic foraminifera had their origin in Tethys. Following the Early Albian migration of the Tethyan foraminifera, however, they seem to have become isolated in the South East Asian sub-province, again correlated with the Early Albian sea-level recovery. During the Late Albian, the lineages evolved independently but in parallel to their Tethyan ancestors. The form *Conicorbitolina* sp. is similar to but different in specific characters from the Tethyan *C. conica* (d'Archiac). This suggests that the migration of Albian foraminifera to the Western Pacific province was only possible for a limited period around the Early Albian. Thereafter the orbitolinids of the South East Asian sub-province remained small, rare and isolated from those in Tethys, as the exclusively Tethyan large species of *Orbitolina* never appeared in this sub-province. The orbitolinids do not survive the Albian-Cenomanian boundary, but unlike the Tethyan realm, they disappeared completely from the Northwest Pacific sub-province at the end of PZ Albian 1 (Iba et al., 2011) and, from the South East Asian sub-province at the end of PZ Albian 4.

CONCLUSION

Analysis of new material combined with a synthesis of the published literature has allowed the understanding of the global evolution and paleobiogeographic distribution of mid-Cretaceous orbitolinids within three LBF provinces; namely the Americas, Tethys and the newly identified Western Pacific province.

We conclude that, unlike previously studied Cenozoic LBF forms, such as the lepidocyclinids (BouDagher-Fadel and Price, 2010), the miogypsinids (BouDagher-Fadel and Price, 2013), the nummulitoids (BouDagher-Fadel and Price, 2014) and the orthophragminids (BouDagher-Fadel and Price, 2017), which evolved first in the Americas and then migrated eastward to Tethys, the Mesozoic orbitolinids originated in the warm tropical shallow platforms of Tethys in the Early Cretaceous, Valanginian (PZ Valanginian 1). The subsequent paleogeographic migration during the global sea-level low stands of the Aptian of members from orbitolinid Group (ii) and Group (v) was bi-directional, moving from Tethys westward to the Americas,

and also eastward into the Western Pacific region. There is no evidence of a West to East trans-Atlantic migration, nor of migration of Western Pacific forms to Tethys.

We infer that migration stopped after rising sea-level in the Albian. As species became geographically isolated, colonizing new but ecologically similar habitats, they thrived and evolved similar but distinct parallel lineages, taking advantages of empty niches and optimum conditions. This example of parallel speciation is discussed by Schluter *et al.* (2004), and probably reflects that all species shared a genetic predisposition to develop mutations of a specific, advantageous type, inherited from their last common ancestor.

The new understanding of the phylogenetic evolution of the Tethyan, Western Pacific and American orbitolinids presented in this paper, when combined with the improved understanding of their biostratigraphic ranges and facies relationships, provides the first global-scale understanding of their development, and so enhances their usefulness as a tool for the study of Early to mid-Cretaceous warm-water carbonate platforms, which are so important in today's hydrocarbon exploration.

Acknowledgements

We would like to thank Prof. Hu Xumian and his students Zhong Han and Wei Zhang of the School of Earth Sciences and Engineering, Nanjing University for their assistance in sampling Tibet sections, and the South East Asia Research Group, Royal Holloway and in particular Dr. Duncan Witts for providing us with Borneo, Kalimantan samples.

REFERENCES

- An, W., Hu, X., Garzanti, E., BouDagher-Fadel, M. K., Wang, J., and Sun, G., 2014. Xigaze forearc basin revisited (South Tibet): Provenance changes and origin of the Xigaze Ophiolite: *Geological Society of America Bulletin*, v. 126, p. 1595–1613.
- Arnaud Vanneau, A., 1975. Réflexion sur le mode de vie de certains Orbitolinidés (Foraminifères) barrémo-aptiens de l'Urgonien du Vercors.: *Compte rendu des séances de la société de physique et d'histoire naturelle de Genève.*, v. 10 fascicule, p. 126-130.

- Arnaud Vanneau, A., 1980. Micropaléontologie, paléoécologie et sédimentologie d'une pl.-forme carbonatée de la marge passive de la Téthys: l'Urgonien du Vercors septentrional et de la Chartreuse (Alpes occidentales): *Géologie Alpine*, v. Mémoire 11, p. 1– 874.
- Banner, F. T., and Simmons, M. D., 1994. Calcareous algae and foraminifera as water-depth indicators: an example from the Early Cretaceous carbonates of Northwest Arabia, *in* Simmons, M. D. (ed.), *Micropalaeontology and Hydrocarbon Exploration in the Middle East*: Chapman and Hall, London, p. 243-252.
- Bosellini, A., 1992. The continental margins of Somalia. Structural evolution and sequence stratigraphy. *American Association of Petroleum Geologists Memoir*, v. 53, p. 185 – 205.
- Bosellini, A., Russo A. , and R. Schroeder 1999. Stratigraphic evidence for an Early Aptian sea-level fluctuation: The Graua Limestone of south-eastern Ethiopia. *Cretaceous Research*, v. 20, p. 783 – 791.
- BouDagher-Fadel, M. K., 2008. Evolution and Geological Significance of Larger Benthic Foraminifera, *Developments in Palaeontology and Stratigraphy*: ELSEVIER, Amsterdam, v. 21, 540 p.
- Boudagher-Fadel, M. K., 2018a. Evolution and Geological Significance of Larger Benthic Foraminifera. London, UK, 704 p.
- Boudagher-Fadel, M. K., 2018b. Revised diagnostic first and last occurrences of Mesozoic and Cenozoic planktonic foraminifera. UCL Office of the Vice-Provost Research, Professional Papers Series, 1–5.

- Boudagher-Fadel, M. K., and Price, G. D., 2013. The phylogenetic and palaeogeographic evolution of the miogypsinid larger benthic foraminifera, 170, 2013, pp. 185–208.
- BouDagher-Fadel, M. K., and Price, G. D., 2014. The phylogenetic and palaeogeographic evolution of the nummulitoid larger benthic foraminifera. *Micropaleontology*, 60, 483–508.
- Boudagher-Fadel, M. K., and Price, G. D., 2017. The paleogeographic evolution of the orthophragminids of the Paleogene. *Journal of Foraminiferal Research*, 47, 337–357.
- Boudagher-Fadel, M. K., Hu, X., Price, G. D., Sun, G., Wang, J. -. G., and An, W., 2017. Foraminiferal biostratigraphy and palaeoenvironmental analysis of the mid-Cretaceous limestones in the southern Tibetan plateau. *Journal of Foraminiferal Research*. 47, no. 2, p. 188–207.
- BouDagher- Fadel, M.K., and Wilson, M., 2000. A revision of some larger Foraminifera of the Miocene of South- East Kalimantan. *Micropaleontology*, 42 (2), 153 – 166.
- Brouwer, H.A., 1934. Geologische onderzoekingen op het eiland Celebes. Kolinie Geologische-Mijnbouw Genootschap, Verhandelingen. Geologische Series 10, 905 39–218.
- Cantrell, D.L., Griffiths, C.M. and Hughes, G.W., 2017. New tools and approaches in carbonate reservoir quality prediction: A case history from the Shu'aiba Formation, Saudi Arabia. In: Agar, S. M. and Geiger, S. (eds) 2015. *Fundamental Controls on Fluid Flow in Carbonates: Current Workflows to Emerging Technologies*. Geological Society, London, Special Publications, 406, 401–424.
- Carter, D.J. and Hart, M.B., 1977. Aspects of mid-Cretaceous stratigraphic micropalaeontology. *Bulletin of the British Museum (Natural History), Geology*, 29, 1-135.

- Caus. E, Teixell, A., and Bernaus, J.M., 1997. Depositional model of a Cenomanian-Turonian extensional basin (Sopeira Basin, NE Spain): interplay between tectonics, Eustasy and biological productivity. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 129, 23-36.
- Cherchi A., Schroeder R. and Bin Ghoth, M., 1998. Early Aptian orbitolinid foraminifera from the Qishn Formation of Al Mukalla (Hadramawt, Southern Yemen). Comparisons with adjacent regions.- *Zeitschrift für geologische Wissenschaften*, Berlin, vol. 26, n° 5/6, p. 543-561.
- Cherchi, A., and Schroeder, R., 1999. Late Barremian orbitolinid Foraminifera from northern Somalia. *Bollettino della Società Paleontologica Italiana*, Modena, 38, 3-13.
- Cherchi, A., and Schroeder, R., 2004. Evolution of orbitolinid foraminifers and anoxic events: A comment on an article by J. Guex, *Eclogae Geologicae Helvetiae*, 97, 441-444.
- Douglass, R. C., 1960. The Foraminiferal Genus *Orbitolina* in North America: Geological Survey Professional Paper, 333, 1–752.
- Görög, A., and Arnaud Vanneau, A., 1996. Lower Cretaceous Orbitolinas from Venezuela: *Micropaleontology*, 42, 65– 78.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D. and Ogg, G. M., 2012. *The Geologic Time Scale 2012*, Elsevier, Amsterdam, 2 volumes, 1144 p.
- Hart, M.B., 1982. The Marine Rocks of the Mesozoic. In: Durrance, E.M. & Lamming, D.J.C. (eds): *The Geology of Devon*, 179-203, Exeter University Press.

- Hart, M.B., and Williams, C.L., 1990. The Upper Greensand in East Devon: new data but old problems. *Proc. Ussher Soc.*, 7, 273-27.
- Hart, M.B., Manley, E.C., and Weaver, P.P.E., 1979. A biometric analysis of an *Orbitolina* fauna from the Cretaceous succession at Wolborough, S. Devon. *Proc. Ussher Soc.*, 4, 317-326.
- Hashimoto, W., and Matsumaru, K., 1974. *Orbitolina* from the Seberuang Cretaceous, Kalimantan Barat (West Borneo), Indonesia. In: T. Kobayashi and R. Toriyama (eds.) *Geology and Palaeontology of Southeast Asia*, Tokyo University Press, 14, p. 89-99.
- Hashimoto, W., and Matsumaru, K., 1977. *Orbitolina* from West Sarawak, east Malaysia. *Geology and Palaeontology of Southeast Asia*, 19: 81-88.
- Hashimoto, W., E. Aliate, N. Aoki, G. Balce et al., 1975. Cretaceous system of Southeast Asia. In: T. Kobayashi and R. Toriyama (eds.) *Geology and Palaeontology of Southeast Asia*, University of Tokyo Press, 145, 219-287.
- Henson, F. R. S., 1948. Larger Imperforate Foraminifera of South-Western Asia (Families Lituolidae, Orbitolinidae and Meandropsinidae): Xi, British Museum (Natural History), London, 127 p.
- Hofker, J., 1963. Studies on the genus *Orbitolina* (Foraminiferida): *Leidse Geologische Medelingen*, v. 29, p. 181-253.
- Hottinger, L., 1978. Comparative anatomy of elementary shell structures in selected larger Foraminifera, in Hedley, R. H., and Adams, C. G. (eds), *Foraminifera 3*: Academic Press, London, p. 203-266.

- Hughes, G. W. 2005. Micropalaeontological dissection of the Shu'aiba reservoir, Saudi Arabia. In: Powell, A. J. & Riding, J. B. (eds). Recent Development in Applied Biostratigraphy. The Micropalaeontological Society, Special Publications, 69–90.
- Husinec, A., 2001. *Palorbitolina lenticularis* from the northern Adriatic region: Palaeogeographical and evolutionary implications: Journal of Foraminiferal Research, 31, 287–293.
- Iba, Y. and Sano, S., 2006. *Mesorbitolina* (Cretaceous larger foraminifera) from the Yezo Group in Hokkaido, Japan and its stratigraphic and paleobiogeographic significance. Proceedings of the Japan Academy Series B, 82, 216-223.
- Iba, Y., Taki, S., Yoshida, K. and Hikida, Y., 2005. *Orbitolina* bearing limestone pebbles from the lowermost part of Lower Yezo Group (Lower Cretaceous) in the Nakagawa area, northern Hokkaido, Japan and its significance. Journal of Geological Society of Japan, 111: 67-73 (in Japanese with English abstract).
- Iba, Y., Sano S.I. and Miura T., 2011. Orbitolinid foraminifers in the Northwest Pacific: their taxonomy and stratigraphy. Micropaleontology, New York, vol. 57, n° 2, p. 163-171.
- Littler, K., Robinson, S.A., Bown, P.R., Nederbragt, A.J., and Pancost, R.D., 2011. High sea-surface temperatures during the Early Cretaceous Epoch, Nature Geoscience, 4, 169-172.
- Luger, P., Hendriks, F., Arush, M., Bussmann, M., Kallenbach, H., Mette, W., and Strouhal, A., 1990. The Jurassic and Cretaceous of northern Somalia: Preliminary results of the sedimentologic and stratigraphic investigations. Berliner geowissenschaftliche Abhandlungen, (A), 120.2, 571-594.

- Masse, J.-P., 1976. Les calcaires urgoniens de Provence (Valanginian-Aptien), Stratigraphie, paléontologie, les paléoenvironnements et leur evolution: Thèse Doctorat D'Etat, Université Aix-Marseille, France, v. I, II, 445 p.
- Masse, J.-P., Chartrousse, A., and Borgomano, J., 1998. The Lower Cretaceous (Upper Barremian-Lower Aptian) Caprinid Rudists from northern Oman. *Géobios, mémoire spécial*, 22, 211-223.
- Matsumaru, K., 1971. Certain larger foraminifera from Japan. *Journal of Saitama University, Faculty of Education*, 20: 149-159.
- Matsumaru, K., 1973. Larger Foraminifera from the Cretaceous and Tertiary in Japan. In, *Atlas of Japanese Fossils Editorial Committee (ed.), Atlas of Japanese Fossils*, 24 pp (in Japanese). Tokyo: Tsukiji Shokan Publishing Co., LTD., 22–127.
- Matsumaru, K., 2005. *Praeorbitolinoides*, a new Orbitolinid foraminiferal genus from the Lower Aptian (Cretaceous) of Hokkaido, Japan. *Micropaleontology*, 51, 93-99.
- Matsumaru, K., and Furusawa, A., 2005. The present condition and subject of the Cretaceous Orbitolinid foraminiferal studies of Japan. *Journal of Saitama University, Faculty of Education*, 54, 15-22 (in Japanese).
- Matsumaru, K., and Furusawa, A., 2007. On Orbitolinid foraminifera from the Lower Aptian (Cretaceous) of Hokkaido, Japan. *Journal of the Palaeontological Society of India*, 52, 39-44.
- Matsumaru, K., Sudo K. and Senaha, T., 1976. A discovery of *Orbitolina* from the calcareous sandstone of the Koma River, Hidaka-cho, Iruma-Gun, Saitama Prefecture, Japan. *Journal of Geological Society of Japan*, 82, 661-662 (in Japanese).

- Maync, W., 1955, *Dictyoconus walnutensis* (Carsey, 1926) in the mid-Albian Guacharo limestone of eastern Venezuela. Contributions from the Cushman Foundation for Foraminiferal Research, 6, part 3, 85-96.
- Mehrnusch, M., 1973. Eine orbitoliniden-Fauna aus der Unterkreide von Esfahan (Zentral Iran). Neues Jahrbuch für Geologie und Paläontologie Monatshefte, Stuttgart, 6, 374-382.
- Meza, J., 1980. El género *Orbitolina* en México y su distribución estratigráfica. Revista del Instituto Mexicano del Petróleo, 12(3), 4-33.
- Miller, K., Kominz, M., Browning, J., Wright, J., Mountain, G., Katz, M., Sugarman, P., Cramer, B., Christie-Blick, N., and Pekar, S., 2005, The Phanerozoic record of global sea-level change. Science, 310, 1293–1298.
- Monreal, R., and Longoria, J.F., 1999. A Revision of the Upper Jurassic and Lower Cretaceous stratigraphic nomenclature of the Chihuahua trough, north central Mexico: Implications for lithocorrelations: In Bartolini, C., Wilson, J.L., Lawton, T.F. (eds.). Geological Society of America Special Paper, 340, 69-92.
- Musper, K.A.F.R., 1937. *Geological map of Sumatra*. Explanatory notes to sheet 16 (Lahat), scale 1:200 000. Dienst Mijnbouw Nederlandsch Indie.
- Naafs, B. D. A., Castro, J. M., De Gea, G. A., Quijano, M. L., Schmidt, D. N., and Pancost, R. D., 2016. Gradual and sustained carbon dioxide release during Aptian Oceanic Anoxic Event 1a. Nature Geoscience, 9, 135–139.

- Omaña, L., and Alencáster, G., 2009. Lower Aptian shallow-water benthic foraminiferal assemblage from the Chilacachapa range in the Guerrero-Morelos Platform, south Mexico. *Revista Mexicana de Ciencias Geológicas*, 26, 3, 575-586.
- Pantoja-Alor, J., Schroeder, R., Cherchi, A., Alencaster, G., and Pons, J.M., 1994. Fossil assemblages, mainly foraminifers and rudists, from the early Aptian of southwestern Mexico. Paleobiogeographical consequences for the Caribbean Region. *Revista Española de Paleontología*, 9, 211-219.
- Peybernès, B., 1982. Les Orbitolinides crétacés d'Afrique; essai de synthèse. In: Blondeau, A., (Editor); Colloque africain de micropaléontologie; IV, Notes consacrées a l'Afrique Occidentale et Equatoriale. Bureau de Recherches géologiques et Minières, Paris, p.13-28. (Cahiers de Micropaléontologie 2).
- Pittet, B., Van Buchem, F. S. P., Hillgärtner, H., Razin, P., Grötsch, J., and Droste, H., 2002. Ecological succession, palaeoenvironmental change, and depositional sequences of Barremian-Aptian shallow-water carbonates in northern Oman. *Sedimentology*, v. 49, p. 555–581.
- Roobahani, P.R., 2011. Lithostratigraphy and biostratigraphy of the Lower Cretaceous of the Jalmajird area (northeast of Khomeyn, Central Iran Basin), Iran. *Geo. Alp*, 9, 48–58.
- Royer D.L., Berner R.A. and Park, J., 2007. Climate sensitivity constrained by CO² concentrations over the past 420 million years. *Nature*, 446, 530-532.
- Saint-Marc, P., 1970. Contribution à la connaissance du Crétacé basal au Liban. *Revue de Micropaléontologie*, 12, 4, 224-233.

- Salnikov, B. A., and Tikhomolov, Y. I., 1987. Stratigraphy, Chapter 3, A description of the section. In: Poyarkova, Z. N. (ed.), Reference section of Cretaceous deposits in Sakhalin (Naiba section), 13-49 (In Russian). Moscow, USSR Academy of Sciences.
- Schluter, D., Clifford, E.A., Nemethy, M. and McKinnon, J.S., 2004. Parallel Evolution and Inheritance of Quantitative Traits. *The American Naturalist*, 163, 809-822.
- Schroeder, R., 1975. General evolutionary trends in orbitolinas. *Revista española de Micropaleontología*, Numero especial, p.117–128.
- Schroeder, R., and Cherchi, A., 1979. Upper Barremian - lowermost Aptian orbitolinid foraminifers from the Grand Banks continental rise, northwestern Atlantic (DSDP Leg 43, Site 384). In B.E. Tucholke, P.R. Vogt et al. (eds.), Initial reports of the Deep Sea Drilling Project, 43, 575-583.
- Schroeder, R., Van Buchem, F. S.P., Cherchi, A., Baghbani, D., Vincent, B., Immenhauser, A., and Granier, B., 2010. Revised orbitolinid biostratigraphic zonation for the Barremian – Aptian of the eastern Arabian Plate and implications for regional stratigraphic correlations. *GeoArabia Special Publication* 4, 1, 49–96.
- Sen Gupta, B. K., and Grant, A. C, 1971. *Orbitolina*, a Cretaceous larger foraminifer, from Flemish Cap: paleoceanographic implications. *Science*, 173, 934-936.
- Shakib, S.S., 1994. Paleoenvironmental and biostratigraphic significance of foraminiferal association from the Early Cretaceous sediments of southwest Iran. - In: Simmons, M.D. (ed.), *Micropalaeontology and Hydrocarbon Exploration in the Middle East*, 127-155, British Micropalaeontological Society Publication Series, Chapman & Hall, London.

- Simmons, M.D., 1994. Micropalaeontological biozonation of the Kahmah Group (Early Cretaceous), central Oman Mountains, *in* Simmons, M.D. (ed.), *Micropalaeontology and Hydrocarbon Exploration in the Middle East*: Chapman & Hall, London, 177–220.
- Simmons, M.D., Whittaker, J.E. and Jones, R.W., 2000. Orbitolinids from Cretaceous sediments of the Middle East—a revision of the F.R.S. Henson and associated collection. In: Hart, M.B., Smart, C.W. (Eds), *Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*: Gybrowski Foundation special publication, 7, 411–437.
- Simmons, M.D., and Hart, M.B., 1987. The biostratigraphy and microfacies of the Early to mid-Cretaceous carbonates of Wadi Mi'aidin, Central Oman Mountains. In M.B. Hart (Ed.), *Micropalaeontology of Carbonate Environments*, p. 176-197.
- Simmons, M.D., Williams, C.L., and Hart, M.B., 1992. Sea-level changes across the Albian-Cenomanian boundary in South-West England. *Proceedings of the Ussher Society*, 7, 408-412.
- Skelton, P.W., and Masse, J.-P., 2000. Synoptic guide to Lower Cretaceous rudist bivalves of Arabia. *Society of Economic Paleontologists and Mineralogists Special Publication*, 69, 89-99.
- Ujiié, H., and Kusukawa, T., 1968. *Orbitolina* (Cretaceous Foraminifera) from the Miyako Group, Iwate Prefecture, Northeast Japan. *Memoirs of the National Science Museum*, Tokyo, 1, 29-33 (in Japanese with English abstract).

- Vahrenkamp, V.C., 1996. Chemostratigraphy on the Lower Cretaceous Shu'aiba formation: a $\delta^{13}C$ reference profile for the Aptian stage from the southern Neo-Tethys Ocean. *American Association of Petroleum Geologists Bulletin*, 80, 647–62.
- Vilas, L., Masse, J.-P., and Arias, C., 1995. *Orbitolina* episodes in carbonate platform evolution, the early Aptian model from SE Spain: Palaeogeography Palaeoclimatology Palaeoecology, 119, 35–45.
- Walliser, O.H., 1995. Global events and event Stratigraphy in the Phanerozoic. In: Walliser, O.H. (ed.), *Global Events and Event Stratigraphy*. Springer, Berlin, 7–19.
- White, L. T., Hall, R., Armstrong, R. A., Barber, A. J., BouDagher-Fadel, M. K., Baxter, A, Wakita, K, Manning, C, and Soesilo, J., 2017. The geological history of the Latimojong region of western Sulawesi. *Journal of Asian Earth Sciences*, 138, 72-91.
- Witt, W., and Gökdağ 1994. Orbitolinid biostratigraphy of the Shu'aiba Formation (Aptian), Oman - implications for reservoir development. In M.D. Simmons (ed.), *Micropalaeontology and Hydrocarbon Exploration in the Middle East*. Chapman and Hall, London, 241-271.
- Witts, D., 2011. *Stratigraphy and Sediment Provenance: The Barito Basin, Southeast Kalimantan, Indonesia*. Unpublished PhD thesis, Royal Holloway University of London.
- Yabe, H., 1946. On some fossils from the Saling Limestone of the Goemai Mts., Palembang, Sumatra- I. *Proc. Japan Academy* 22, 6, 200-203.
- Yabe, H., and Hanzawa, S., 1926. Geological age of *Orbitolina* bearing rocks of Japan. *Science Reports of Tohoku Imperial University, Second Series*, 9: 13-20.

PLATE CAPTIONS

Plate 1

Scale bars: Figs 1, 3 = 1mm; Figs 2, 4-8 = 0.5mm. Key words: marg = marginal zone; rad = radial zone; ret = reticular zone. All samples are deposited in School of Earth Sciences and Engineering Nanjing University.

Figs 1-3, 6. *Mesorbitolina aperta* (Erman), Langsham Formation, Tibet, PZ Albian 3-Cenomanian 1. 1) Axial section through the megalospheric embryonic apparatus; 2) basal section of the megalospheric embryonic apparatus; 3) thin section showing the details of the radial zone; 6) thin section through the reticular zone.

Fig. 4. *Mesorbitolina* sp., Tibet, Aptian. Thin section through the marginal zone.

Fig. 5. *Mesorbitolina subconcava* Leymerie, Indonesia, PZ Albian 1. Basal section through embryonic apparatus showing the periembryonic chambers.

Figs 7-8. *Mesorbitolina texana* (Roemer), Tibet, PZ Aptian 4. Random thin sections: 7) showing details of the radial and reticular zones; 8) basal view showing the zigzag main partitions with apertural pores at the reentrants. The partitions are broken up in the central complex reticular zone.

Plate 2

Scale bars = 1mm. All samples are deposited in School of Earth Sciences and Engineering Nanjing University.

Fig. 1. *Palorbitolina lenticularis* (Blumenbach), Jiarong section, TLK1a, PZ Aptian 2, sample 14LZ13.

Fig. 2. *Praeorbitolina cormyi* Schroeder, Laxue section, TLK1a, PZ Aptian 2, 14 LZ12

Fig. 3. *Palorbitolinoides orbiculata* Zhang, Langsham section, TLK1a, PZ Aptian 2, 16SL 02.

Fig. 4. *Mesorbitolina aperta* (Erman). Guolong section, TLK1h, PZ Cenomanian 1, 13GL33.

Fig. 5. *Palorbitolinoides hedinii* Cherchi and Schroeder, Langsham section, TLK1e, PZ Albian 2, 16SL45.

Fig. 6. *Mesorbitolina subconcava* Leymerie, Langsham section, TLK1c, PZ Aptian 4b, 16SL29.

Plate 3

Scale bars = 1mm. All photos are from sections from the western flank of the Meratus Mountains, Barito Basin, Southeast Kalimantan, Indonesia, All samples are deposited in UCL Collections.

Figs 1-3. *Mesorbitolina texana* (Roemer), PZ Albian 1, BBr-14. 2-3) vertical sections.

Figs 4, 7. *Palorbitolinoides orbiculata* Zhang, PZ Albian 1, BBr-14.

Fig. 5. *Mesorbitolina subconcava* Leymerie, PZ Albian 1, BBr-14, oblique transverse section through embryonic apparatus.

Fig. 6. *Conicorbitolina* sp., PZ Albian 4, BBr-22.

FIGURE CAPTIONS

Figure 1. The diagnostic first and last occurrences of Cretaceous planktonic foraminiferal species, calibrated against the most recent biostratigraphic time scale and radio-isotope data (after BouDagher-Fadel, 2018b).

Figure 2. (A) Simplified tectonic map of the Tibetan Plateau and adjacent regions, showing the Lhasa terrane in the context of the Tibetan Plateau (Pan et al., 2004). JSSZ—Jinsha suture zone; BNSZ—Bangong-Nujiang suture zone; IYSZ—Indus-Yarlung suture zone. (B) Simplified geological map of the Lhasa terrane modified from Kapp et al. (2005). SGAT—Shiquan-Gaize-Amdo thrust; GST—Gaize–Selin Co thrust; GLT—Gugu La thrust; ST—Shibalu thrust; ELT—Emei La thrust; GT—Gangdese thrust system; GCT—Great Counter thrust. Section 1 from the Xigaze forearc basin; Sections 2 and 3 from the Linzhou basin; Sections 4, 5 and 6 from the Coqen basin.

Figure 3. Cenozoic geology of the Barito and Asem-Asem Basins (modified from Supriatna et al., 1994; Witts, 2011).

Figure 4. Gradual morphological changes from primitive orbitolinids to the advanced *Orbitolina* in Tethys.

Figure 5. Example of evolutionary Tethyan lineages from Group (ii) to (v).

Figure 6. Variation in sea-level during the mid-Cretaceous based on Miller et al. (2005) correlated to the boundaries of the PZ after BouDagher-Fadel (2018b) and showing dominant assemblages at the top of regression and transgression phases.

Figure 7. The provincial distribution of the orbitolinids during Early Cretaceous in the Tethys (1), the Western Pacific (2), and the Americas (3), with paleo-oceanic currents shown by the white arrows.

Figure 8. The test architecture of *Orbitolina* (not to scale). (1) Test dissected in several places to show the internal structures (after Douglass, 1960); (2) Diagrams showing micro-structures of *Orbitolina* exposed by tangential sections cut progressively deeper below the epidermis:

- a- Megalospheric embryonic apparatus, b- Slightly eroded surface exposing sub-epidermal cells; b1/b2- regular/irregular arrangement of secondary epidermal cells (stage III),

- c- Primary sub-epidermal cells (stage II),
- d- Marginal chamberlets (stage I) with residual traces of vertical primary sub-epidermal plates only, d1-d4- Sections through marginal chamberlets between (c and d) and the beginning of the true radial chamber-passages with canals (e1, e3, e5, e7).
- e1-e2- Radial chamber passages sub-rounded, canals short, wall thickness relatively small,
- e3-e4- Radial chamber passages triangular, canals long, wall thickness relatively great,
- e5-e6- Radial chamber passages initially rectangular, with simple perforations, wall thickness small, e7-e8- Radial chamber passages irregular – rounded originating from vertical pairs of primary sub-epidermal cells, canals short, wall thickness small, f- Main triangular partitions with zigzag shape when seen deeper in the test, g- the complex central zone.

Figure 9, (a-b) – Diagrammatic axial sections of orbitolinids, a- showing closely spaced chamber layers with primary horizontal sub-epidermal plates only; b- showing widely-spaced chamber layers with primary and secondary horizontal sub-epidermal plates. (c-e) Diagrammatic basal sections of orbitolinids, c- marginal zone broad, marginal chamberlets triangular, radial walls thick, straight; d- marginal zone broad, marginal chamberlets rectangular, radial walls thin, zigzag; e- marginal zone narrow, marginal chamberlets sub-triangular, radial walls moderately thick, vertical primary sub-epidermal plates thickening inward with some prolonged as radial walls. Note: In species having triangular radial passages, the thickness of radial walls as seen in basal views will vary according to the position on the section just above or just below a chamber floor. (f-j) Diagrams illustrating radial and reticular zones of orbitolinids as seen in basal views; f- radial partitions; g- reticular partitions, h- radial chamber passages, i- radial and reticular chamber passages, j- complex reticular zone, no radial zone. (k-r) Diagrams showing various textures of the central zone in *Orbitolina* as observed in axial and oblique sections; shaded areas and lines represent shell material. (k-o) axial sections; k – Wall and floor thickness and chamber diameters sub-equal; chamber layers clearly marked and connected by short, sub-vertical canals; l – wall and floors thin; chamber layers clearly connected by simple perforations and clearly recognisable by alignment of longitudinal segments of chamber passages in the radial zone; m – wall and floors thin; chamber layers clearly connected by simple perforations but not clearly recognisable; the section is cut through the reticular zone and chamber segments are all more or less transverse;

n – wall thickness small, floor thickness relatively great; chamber layers clearly marked and connected by oblique canals not visible in axial sections; o, wall and floor thickness s great; chamber layers not clearly recognizable owing to wide spacing of chamber segments; oblique canals not visible in axial sections. (p-r) oblique sections; p – Oblique section corresponding to (a) above; labyrinthic texture; canals (when visible) not clearly differentiated from chamber segments; q – Oblique section corresponding to (c) above; that corresponding to (b) would be similar but would show a few longitudinal chamber segments; cellular texture; r – Oblique section corresponding to (d) or (e) above; dendritic texture; small chamber segments connected by long, oblique canals forming a roughly polygonal network.

Figure 10, Enlargement of parts of *Palorbitolina lenticularis* (Blumenbach) figured by BouDagher-Fadel (2018a), scale bar = 100 μm . (A) Transverse section showing the triangular main partitions; (B) the same transverse section showing the central complex reticular part of the test; (C) the same transverse section showing the subdivision of the marginal chamberlets into cellules/chamberlets.

Figure 11. Range chart for some key orbitolinid species from Group (v) in the Tethyan and Western Pacific, and American provinces.

Figure 12. The facies range of the dominant orbitolinids in a Tethyan carbonate shelf. Integrated reef/ramp model for Neogene carbonates. The ramp model is indicated by the blue dotted line. In the case of gently sloping ramp, the outer ramp lithofacies are made of mudstones and wackestones, while in the middle ramp mudstone with carbonate nodules would develop.

Plate 1










Plate 2








Plate 3






Fig. 1

Age (Ma)	Period, Epoch, Stage	Planktonic zonation	Diagnostic First Occurrence of Planktonic Foraminifera		Diagnostic Last Occurrence of Planktonic Foraminifera				
			Age (Ma)	Species	Age (Ma)	Species			
66.0	Cretaceous	Late	3	b	66.2	<i>Ventilabrella multicamerata</i>	66.0	<i>Globotruncana</i> spp./ <i>Rugoglobigerina</i> spp.	
				a	67.0	<i>Plummerita reicheli</i>	66.2	<i>Contusotruncana plicata</i>	
			2		70.0	<i>Racemiguembelina powelli</i>	67.0	<i>Rugotruncana subpennyi</i>	
				1	b	71.0	<i>Contusotruncana plicata</i>	70.0	<i>Globotruncana lapparenti</i>
			a		72.1	<i>Plummerita hantkeninoides</i>			
			Camp.	3	b	74.5	<i>Rugotruncana subpennyi</i>	72.1	<i>Radotruncana subspinoso</i>
					a	77.0	<i>Globotruncanita stuarti</i>	74.5	<i>Globotruncanita atlantica</i>
						80.3	<i>Radotruncana calcarata</i> , <i>Globotruncanita atlantica</i>	77.0	<i>Radotruncana calcarata</i>
			Sant.	2		83.6	<i>Globotruncana mariei</i>	80.3	<i>Rugoglobigerina subrugosa</i>
					1		84.9	<i>Rugoglobigerina subrugosa</i>	83.6
			b	85.6		<i>Globotruncana linneiana</i>			
			Con.	2		86.3	<i>Concavatotruncana asymetrica</i>	86.3	<i>Marginotruncana paraconcovata</i>
						88.0	<i>Globotruncana lapparenti</i>		
			Tur.	3		89.8	<i>Marginotruncana paraconcovata</i>	89.8	<i>Sigalitruncana biconvexiformis</i>
						91.3	<i>Dicarinella primitiva</i>	91.3	<i>Helvetoglobotruncana helveticae</i>
			Cen.	2		92.8	<i>Helvetoglobotruncana helveticae</i>	92.8	<i>Concavatotruncana elata</i>
						93.9	<i>Sigalitruncana biconvexiformis</i>	93.9	<i>Rotalipora cushmani</i>
			Alb.	4	b	95.1	<i>Rotalipora cushmani</i>	95.1	<i>Favusella washitensis</i>
					a	97.2	<i>Thalmaninella reicheli</i>		
			Apt.	3	b	98.8	<i>Thalmaninella greenhornensis</i>	99.6	<i>Alanlordella bentonensis</i>
					a	100.5	<i>Thalmaninella globotruncanoides</i>		
			Barr.	2	b	101.5	<i>Thalmaninella gandolfi</i>	100.5	<i>Hedbergella rischi</i>
					a	102.2	<i>Ticinella madecassiana</i>		
			Haut.	3	b	104.4	<i>Pseudothalmaninella subticinensis</i>	102.2	<i>Ticinella praeticinensis</i>
					a	106.7	<i>Ticinella praeticinensis</i>		
			Val.	2		109.8	<i>Ticinella roberti</i>	106.7	<i>Blefuscuiana aptiana</i>
	111.4	<i>Ticinella primula</i>							
Berr.	1	b	113.0	<i>Hedbergella rischi</i>	113.0	<i>B. maslakovae/ L. globulifera</i>			
		a	115.0	<i>Blefuscuiana kuznetsovae</i>					
Tith.	4	b	116.5	<i>Lilliputianella kuhryi</i>	119.5	<i>Praehedbergella ruka</i>			
		a	119.5	<i>Praehedbergella tatianae</i>					
Jurassic	Tith.		122.9	<i>Blefuscuiana infracretacea</i>	122.9	<i>Gorbachikella anteroapertura</i>			
			125.0	<i>Blefuscuiana maslakovae</i>	125.0	<i>Gorbachikella grandiapertura</i>			
	Barr.	3		127.0	<i>Lilliputianella globulifera</i>	129.0	<i>Praehedbergella tuschepsensis</i>		
				129.0	<i>Blefuscuiana daminiae</i>				
	Haut.	2		129.4	<i>B. aptiana/Pr. tuschepsensis</i>	132.4	<i>Praehedbergella handousi</i>		
				132.4	<i>Gorbachikella anteroapertura</i>				
	Val.	2		132.9	<i>Praehedbergella handousi</i>	136.65	<i>Conoglobigerina gulekhensis</i>		
				136.6	<i>Gorbachikella grandiapertura</i>				
	Berr.	1		139.8	<i>Gorbachikella sp.</i>	139.8	<i>Conoglobigerina sp. A</i>		
				145.0	<i>Conoglobigerina gulekhensis</i>				
	Tith.			145.0	<i>Conoglobigerina conica</i>	145.0	<i>Conoglobigerina conica</i>		
				152.1	<i>Conoglobigerina conica</i>				

Fig. 2




Fig. 3




Fig. 4




Fig. 6




Fig. 7




Fig. 8





Fig. 9







Fig. 10



A



B



C

Fig. 11




Fig. 12

